Purpose of the Report

The *Dialysis Facility Report (DFR) for Fiscal Year (FY) 2026* is provided as a resource for characterizing selected aspects of clinical experience at this facility relative to other dialysis facilities in this state, ESRD Network, and across the United States. Since these data could be useful in quality improvement and assurance activities, each state's surveying agency may utilize this report as a resource during the FY 2026 survey and certification process.

This report has been prepared for this facility by the University of Michigan Kidney Epidemiology and Cost Center (UM-KECC) with funding from the Centers for Medicare & Medicaid Services (CMS) and is based primarily on data reported in End Stage Renal Disease Quality Reporting System (EQRS), Medicare claims and data collected for CMS. It is the thirtieth in a series of annual reports. This is one of 8,241 reports that have been distributed to ESRD providers in the U.S.

This DFR includes data specific to CCN(s): 999999

Overview

This report includes summaries of patient characteristics, treatment patterns, and patient outcomes for chronic dialysis patients who were treated in this facility between January 2021 and December 2024. Mortality, hospitalization, transplantation, and waitlist statistics are reported for a three- or four-year period. Regional and national averages are included to allow for comparisons. Some of these summaries are adjusted to account for the characteristics of the patient mix at this facility, such as age, sex and diabetes as a cause of ESRD. Unless otherwise specified, summaries include all dialysis patients.

Selected facility highlights from this report are provided on pages 2 through 5 and key data elements that are updated quarterly appear in bold font. For a complete description of the methods used to calculate the statistics in this report, please see the *Guide to the Dialysis Facility Reports for FY 2026* available for download from the methodology section of the Dialysis Data website at www.DialysisData.org.

What's New This Year

The following change has been incorporated into the DFR for FY 2026: the prevalent Standardized Waitlist Ratio (PSWR) was added to Table 6. This measure is similar to the Standardized Waitlist Ratio in the First Year (FySWR), but includes all eligible patients and includes time to all waitlists as opposed to just the first event. The PSWR improves reporting of access to transplantation for prevalent patients compared to the age-adjusted percentage of patient-months waitlisted reported in row 6e, given the focus on time to waitlisting (as opposed to percentage waitlisted in a year), and also includes more robust risk adjustments (similar to the FySWR with the inclusion of patient vintage).

How to Submit Comments

Between August 15, 2025 and August 31, 2025, facilities may submit comments to their state surveyor or UM-KECC by visiting www.DialysisData.org, logging on to view their report, and clicking on the **Comments & Inquiries** tab. Questions or comments after the comment period is over may be submitted to us directly at DialysisData@umich.edu.

- (1) **State Surveyor:** Select **"DFR: Comments on DFR for State Surveyor"** from the drop down list to submit comments regarding this report for the state's surveyor(s). Any comments submitted will be appended and sent to the state's surveyor(s) in September 2025. Please do not include questions for UM-KECC using this option.
- (2) **UM-KECC:** Select **"DFR: Comments on DFR for UM-KECC"** to submit questions or suggestions to improve the DFR to UM-KECC. These comments will not be shared with your state surveyor.

Facility Highlights

Bar charts in this section are displayed as a percentage for all measures reported and highlight the facility's value compared to the state, network, and US.

The line charts in this section are displayed for all standardized measures. The markers show the values of the corresponding standardized measure for this facility, state, network, and US. The bolded horizontal line shows the range of uncertainty due to random variation (95% confidence interval; significant if it does not cross the 1.0 reference line). Regional and national values are plotted above the dotted line to allow for comparisons to facility values.

Patient Characteristics (Tables 1 and 2):

- Among the 15 incident patients with Medical Evidence Forms (CMS-2728) indicating treatment at this facility during 2024:
 - 15% of these patients were not under the care of a nephrologist before starting dialysis, compared to 15% in your State, 15% in your Network, and 16% nationally.
 - •93% of these patients were informed of their transplant options, compared to 94% in your State, 94% in your Network, and 94% nationally.
- Among the patients treated at this facility on December 31, 2024, 14% were treated in a nursing home during the year, compared to 14% in your State, 14% in your Network, and 13% nationally.



Standardized Mortality Ratio (SMR) (Table 3):

- At this facility, the 2021-2024 SMR is 0.98, which is 2% fewer deaths than expected at this facility. Among all U.S. facilities, 50% of facilities had a four-year SMR lower than 0.98. This difference is not statistically significant (p>=0.05), so this lower mortality could plausibly be just a chance occurrence. The 2021-2024 SMR of observed to expected deaths is 0.97 and 0.97 for your State and Network, respectively.
- At this facility, the 2021-2023 first-year SMR of observed to expected deaths is 1.01, which is 48% more deaths than expected at this facility. Among all U.S. facilities, 51% of facilities had a first-year SMR lower than 1.01. This difference is not statistically significant (p>=0.05), so this higher mortality could plausibly be just a chance occurrence. The first-year SMR (2021-2023) of observed to expected deaths is 1.01 and 1.01 for your State and Network, respectively.

2021-2024 SMR

2021-2023 First-Year SMR



Hospitalizations and Readmissions (Table 4):

- The 2021-2024 Standardized Hospitalization Ratio (SHR-Admissions) at this facility is 0.98, which is 2% fewer admissions hospitalized than expected. This difference is not statistically significant (p>=0.05), so this lower hospitalization could plausibly be just a chance occurrence. The 2021-2024 SHR (Admissions) for your State and Network is 0.98 and 0.98, respectively.
- The 2024 Standardized Readmission Ratio (SRR) at this facility is 0.99, which is 1% fewer admissions than expected. This difference is not statistically significant (p>=0.05), so this lower number of readmissions could plausibly be just a chance occurrence. The 2024 SRR for your State and Network is 1.02 and 1.03, respectively.



Infection (Tables 4 and 11):

- The percentage of Medicare dialysis patients at this facility hospitalized with septicemia during 2021-2024 is 12%, compared to 13% in your State, 13% in your Network, and 12% nationally.
- The 2024 rate of PD catheter-related infection was 2.6 per 100 PD patient-months, compared to 2.4 in your State, 2.4 in your Network, and 2.4 nationally.





Transplantation (Table 5):

• The 2021-2024 Standardized 1st Transplantation Ratio (STR) of observed to expected number of patients transplanted for this facility is 1.04, which is 4% higher than expected for this facility. This difference is not statistically significant (p>=0.05) and is plausibly due to random chance. The 2021-2024 STR for your State and Network is 1.05 and 1.04, respectively.

Transplant Waitlist (Table 6):

- The 2024 age-adjusted percent waitlisted at this facility is 16.2%, which equal to the national adjusted percentage. The age-adjusted percent waitlisted in your State and Network is 16.2% and 16.2%, respectively.
- At this facility, the 2021-2023 first-year Standardized Waitlist Ratio (FySWR) is 0.98, which is 2% fewer patients on the waitlist and living donor transplants than expected at this facility. This difference is not statistically significant (p>=0.05) and could plausibly be due to a chance occurrence. The 2021-2023 FySWR for your State and Network is 1.04 and 1.03, respectively.



Influenza Vaccination (Table 7):

• Among the 61 dialysis patients treated at this facility on December 31, 2024, 77% were vaccinated between August 1 and December 31, 2024 compared to 78% nationally. This difference is not statistically significant (p>=0.05) and is plausibly due to random chance. The percentage of patients vaccinated in your State and Network is 78% and 78%, respectively.

Anemia Management (Table 8):

- In 2024, 23% of eligible hemodialysis patient-months had a hemoglobin value below 10 g/dL, compared to 23% in your State, 23% in your Network, and 23% nationally.
- In 2024, 22% of eligible peritoneal dialysis patient-months had a hemoglobin value below 10 g/dL, compared to 22% in your State, 22% in your Network, and 23% nationally.





Dialysis Adequacy (Table 9):

- In 2024, 95% of eligible hemodialysis patient-months had a Kt/V >=1.2 reported, compared to 97% in your State, 97% in your Network, and 97% nationally.
- In 2024, 90% of eligible peritoneal dialysis patient-months had a Kt/V >=1.7 reported, compared to 92% in your State, 92% in your Network, and 92% nationally.

Mineral Metabolism (Table 10):

- In 2024, 17% of eligible patient-months had a serum phosphorus value >7.0 mg/dL, compared to 17% in your State, 17% in your Network, and 17% nationally.
- In 2024, 1.3% of eligible patient-months had calcium uncorrected value >10.2 mg/dL, compared to 1.2% in your State, 1.2% in your Network, and 1.1% nationally.

Vascular Access (Table 11):

- This facility's 2024 Standardized Fistula Rate (SFR) for prevalent patients is 58%, which is 0% higher than the national SFR. This difference is not statistically significant (p>=0.05) and could plausibly be due to a chance occurrence. The SFR in your State and Network is 58% and 58%, respectively.
- Of the prevalent patients receiving hemodialysis treatment at this facility in 2024, the long-term catheter rate was 19%, compared to 18% in your State, 18% in your Network, and 18% nationally.



Dialysis Facility Report for Fiscal Year 2026 SAMPLE Dialysis Facility State: XX Network: 99 CCN: SAMPLE

The tables below provide updates of COVID-19 patient counts, deaths, and hospitalizations among Medicare dialysis patients (Table C1) and Medicare nursing home (NH) dialysis patients (Table C2), 2021-2024. State, Network, and National averages for 2024 are reported for comparison.

TABLE C1. COVID among Medicare Dialysis Patients, 2021-2024^{*1}

			This I	Facility		Regio	onal Averages	, 2024
	Measure Name	2021	2022	2023	2024	State	Network	U.S.
1	Medicare dialysis patients (n)	68	82	71	69	n/a	n/a	n/a
2	Patients ever infected with COVID (n)	11	20	19	12	n/a	n/a	n/a
	Patients ever infected with COVID (% of 1)	16.2	24.4	26.8	17.4	21.7	20.8	22.8
3	Patients initially infected with COVID (n)	3	5	6	0	n/a	n/a	n/a
	Patients initially infected with COVID (% of 1)	4.4	6.1	8.5	0.0	3.7	3.4	3.4
	Deaths among Medicare patients							
4	Deaths (n)	9.0	11.0	13.0	6.0	n/a	n/a	n/a
5	Deaths among patients ever infected with COVID (n)	1.0	3.0	8.0	2.0	n/a	n/a	n/a
	Deaths among patients ever infected with COVID (% of 4)	11.1	27.3	61.5	33.3	30.0	28.3	30.6
	Hospitalizations among Medicare patients *2							
6	Hospitalizations (n)	38	42	39	30	n/a	n/a	n/a
7	Hospitalizations among patients ever infected with COVID (n)	6	10	11	7	n/a	n/a	n/a
	Hospitalizations among patients ever infected with COVID (% of 6)	15.8	23.8	28.2	23.3	24.7	24.3	26.4

TABLE C2. COVID among Medicare Dialysis Patients Treated in Nursing Home (NH) Facilities, 2021-2024^{*1}

		This F	Facility		Regional Averages, 2024			
Measure Name	2021	2022	2023	2024	State	Network	U.S	
Medicare NH dialysis patients (n)	13	19	23	16	n/a	n/a	n/a	
Patients ever infected with COVID (n)	3	5	6	3	n/a	n/a	n/a	
Patients ever infected with COVID (% of 1)	23.1	26.3	26.1	18.8	29.6	28.6	31.7	
Patients initially infected with COVID (n)	1	2	2	0	n/a	n/a	n/a	
Patients initially infected with COVID (% of 1)	7.7	10.5	8.7	0.0	6.2	5.7	6.4	
Deaths among Medicare NH patients								
Deaths (n)	4	6	7	4	n/a	n/a	n/a	
Deaths among patients ever infected with COVID (n)	0	2	4	1	n/a	n/a	n/a	
Deaths among patients ever infected with COVID (% of 4)	0.0	33.3	57.1	25.0	36.7	32.7	36.1	
Hospitalizations among Medicare NH patients *2								
Hospitalizations (n)	10	14	18	14	n/a	n/a	n/a	
Hospitalizations among patients ever infected with COVID (n)	2	3	5	2	n/a	n/a	n/a	
Hospitalizations among patients ever infected with COVID (% of 6)	20.0	21.4	27.8	14.3	29.5	29.5	32.0	

n/a = not applicable

[*1] See *Guide, Section XIX* for detailed information regarding information reported in Tables C1 and C2. Includes patients with Medicare as primary insurer or with a Medicare Advantage plan. Table C2 includes patients who were treated in a nursing home at least one day during the year.[*2] Hospitalization is defined as having at least one day in a hospital from Medicare inpatient claims during the reporting period.

TABLE 1: Summaries for All Dialysis Patients Treated as of December 31st of Each Year, 2021-2024^{*1}

			This F	acility		Region	al Averages*	^A , 202
	Measure Name	2021	2022	2023	2024	State	Network	U.S
a	Patients treated on 12/31 (n)	59	58	58	58	61.7	61.8	61.9
lb	Average age (years)	62.1	62.2	62.3	62.5	62.6	62.7	62.0
c	Age (% of 1a; sums to 100%)							
	< 18	0.9	0.9	0.9	0.9	0.2	0.2	0.2
	18-64	50.8	50.5	49.9	49.3	50.4	50.4	50.
	65+	48.3	48.7	49.2	49.8	49.4	49.3	49.
d	Female (% of 1a)	42.2	41.9	41.8	41.6	41.9	41.9	41.
e	Race (% of 1a; sums to 100%)							
	African American	33.8	33.9	33.7	33.7	35.9	36.1	34.
	Asian/Pacific Islander	5.4	5.6	5.7	5.7	5.9	6.0	7.
	Native American	1.4	1.4	1.5	1.5	1.4	1.3	1.
	White	59.0	58.5	58.1	57.3	55.2	54.9	55
	Other/Unknown/Missing	0.4	0.7	1.1	1.8	1.7	1.7	1.
f	Ethnicity (% of 1a; sums to 100%)							
	Hispanic	18.6	19.3	19.9	20.4	20.8	21.1	24
	Non-Hispanic	81.4	80.7	80.0	79.5	79.1	78.9	75
	Unknown	0.0	0.0	0.1	0.1	0.0	0.0	0.
g	Primary Cause of ESRD (% of 1a; sums to 100%)							
-	Diabetes	44.6	44.0	43.6	42.9	43.5	43.6	44
	Hypertension	30.2	30.4	30.6	30.9	31.6	31.6	31
	Glomerulonephritis	9.5	9.2	8.9	8.7	8.6	8.6	8.
	Other/Unknown	15.4	16.0	16.5	17.0	16.0	15.9	15
	Missing	0.3	0.4	0.4	0.5	0.3	0.3	0.
h	Average duration of ESRD (years)	4.8	4.8	4.8	4.9	5.0	5.0	5.
i	Years since start of ESRD (% of 1a; sums to 100%)							
	<1	17.8	17.3	17.2	17.0	16.5	16.5	16
	1-2	17.2	18.4	17.9	17.7	17.5	17.5	17
	2-3	14.0	13.5	14.4	14.0	13.9	13.9	13
	3-6	25.4	25.3	25.3	25.9	26.1	26.1	26
	6+	25.7	25.5	25.3	25.4	26.1	26.1	26
j	Nursing home patients $(\% \text{ of } 1a)^{*2}$	13.4	13.3	13.5	13.7	13.6	13.6	13
-	Modality (% of 1a; sums to 100%)							
	In-center hemodialysis	84.0	83.3	82.9	82.8	83.0	83.0	83
	Home hemodialysis	2.9	3.2	3.4	3.6	4.1	4.1	3.
	Continuous ambulatory peritoneal dialysis	1.5	1.5	1.6	1.7	1.5	1.5	1.
	Continuous cycling peritoneal dialysis	11.3	11.7	11.9	11.5	11.1	11.1	11
	Other modality	0.4	0.4	0.3	0.3	0.3	0.3	0.

n/a = not applicable

[*A - *F] See Appendix: Table Footnotes on page 29.

[*1] See *Guide, Section IV* for detailed information regarding the summaries included in Table 1.

[*2] Includes patients who were also treated by a nursing facility at any time during the year according to the Nursing Home Minimum Dataset.

SAMPLE Dialysis Facility State: XX Network: 99 CCN: SAMPLE

TABLE 2: Characteristics of New Dialysis Patients, 2021-2024 (Form CMS-2728) *1

			This F	facility		Regional Averages *A			
	Measure Name	2021	2022	2023	2024	State	Network	U.S.	
	Patient Characteristics						2024		
2a	Total number of patients with forms (n)	15	15	15	15	16.1	16.1	16.2	
2b	Average age (years [0-95]) ^{*2}	62.7	62.6	62.8	62.8	63.0	63.0	63.0	
2c	Female (% of 2a)	41.9	41.4	41.4	41.5	41.6	41.6	41.5	
2d	Race (% of 2a; sums to 100%)								
	African-American	29.0	29.4	28.7	28.6	29.1	28.9	27.8	
	Asian/Pacific Islander	5.4	5.6	5.6	5.6	5.7	5.7	6.4	
	Native American	1.3	1.3	1.4	1.4	1.2	1.2	1.2	
	White	63.5	62.2	62.1	59.8	59.6	59.8	59.9	
	Other/Unknown	0.7	1.5	2.2	4.6	4.4	4.4	4.8	
2e	Hispanic (% of 2a)	17.4	17.8	18.7	19.1	19.2	19.4	21.6	
2f	Primary cause of ESRD (% of 2a; sums to 100%)								
	Diabetes	45.0	43.7	43.2	42,4	42.8	42.9	43.4	
	Hypertension	29.7	30.7	31,1	31.3	32.0	32.0	31.7	
	Primary glomerulonephritis	6.0	6.1	5.9	5.8	5.7	5.7	5.7	
	Other/Unknown	19.3	19.5	19.7	20.5	19.4	19.4	19.2	
2g	Medical coverage (% of 2a; sums to 100%)	1910		1717				17.2	
-8	Employer group only	11.7	11.6	11.2	11.5	11.6	11.5	11.5	
	Medicare only	44.7	45.6	45.4	41.5	41.2	41.2	40.6	
	Medicaid only	14.9	15.0	14.9	14.7	14.6	14.7	15.3	
	Medicare and Medicaid only	6.6	6.7	7.5	8.6	8.9	8.9	9.1	
	Medicare and other	7.8	8.1	8.2	9.8	9.8	9.9	9.6	
	Other/Unknown	0.5	9.0	8.9	9.9	9.8	9.8	10.0	
	None	4.9	4.1	3.8	4.1	4.0	4.0	4.0	
2h	Median body mass index (Median; Weight/Height^2)	π.2	7.1	5.0	4.1	4.0	4.0	4.0	
211	Male	28.5	28.4	28.1	28.0	27.7	27.7	27.6	
	Female	29.9	20.4 29.4	29.1	28.0 29.0	28.5	28.5	27.0	
2i	Employment (% of 18-60 year olds in 2a)	29.9	29.4	29.1	29.0	28.5	20.5	20.4	
21	Six months prior to ESRD treatment	33.4	36.2	36.2	36.4	36.8	36.6	36.8	
	At first ESRD treatment	23.9	26.8	26.7	25.5	26.0	25.9	25.9	
2:	Primary modality (% of 2a; sums to 100%)	23.9	20.8	20.7	23.3	20.0	23.9	23.9	
2ј	Hemodialysis	87.2	86.2	85.8	86.5	86.1	86.0	86.1	
	CAPD/CCPD	12.8	13.8	14.2	13.5	13.9	14.0	13.9	
	Other/Unknown	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
2k	Number of incident hemodialysis patients (n)	13	13	13	13	13.9	13.9	13.9	
		15	15	15	15	13.9	13.9	13.9	
21	Access used at first outpatient dialysis (% of 2k; sums to 100%) Arteriovenous fistula	12.0	12.2	14.0	12.6	12.0	12.0	12.0	
		13.0	13.3	14.0	13.6	12.9	12.9	13.0	
	Arteriovenous graft	2.5	2.9	3.2	3.3	3.1	3.1	3.0	
	Catheter	84.2	83.5	82.4	82.6	83.6	83.5	83.5	
2	Other/Unknown/Missing	0.3	0.4	0.4	0.6	0.5	0.5	0.5	
2m	Arteriovenous fistula placed (% of 2k)	23.9	23.2	23.9	22.5	21.4	21.4	21.6	
	Average Lab Values Prior to Dialysis *2								
2n	Hemoglobin (g/dL [5-20])	9.4	9.3	9.3	9.4	9.4	9.4	9.4	
2o	Serum albumin (g/dL [0.8-6.0])	3.4	3.4	3.4	3.5	3.5	3.5	3.5	
2p	Serum creatinine (mg/dL [0-33])	6.5	6.6	6.5	6.4	6.4	6.4	6.4	
	GFR (mL/min [0-30])	9.9	9.8	9.9	10.0	10.1	10.1	10.0	

SAMPLE Dialysis Facility State: XX Network: 99 CCN: SAMPLE

TABLE 2 (cont.): Characteristics of New Dialysis Patients, 2021-2024 (Form CMS-2728)

			This F	acility		Regional Averages *A			
	Measure Name	2021	2022	2023	2024	State	Network	U.S.	
	Care Prior to ESRD Therapy								
2r	Received ESA prior to ESRD (% of 2a)	15.5	16.1	16.5	17.4	17.3	17.4	17.4	
2s	Pre-ESRD nephrologist care (% of 2a;sums to 100%) ^{*B}	15.5	10.1	10.5	17.4	17.5	17.4	17.4	
23	No	17.1	16.5	15.5	15.5	15.3	15.3	15.7	
	Yes, < 6 months	18.5	19.0	19.5	19.1	18.9	19.0	19.0	
	Yes, 6-12 months	18.7	19.0	19.4	19.1	18.5	19.6	19.0	
	Yes, > 12 months	29.1	28.6	29.1	29.4	29.5	29.4	29.0	
	Unknown/Missing	16.5	17.2	17.2	17.0	17.6	17.6	17.6	
2t	Informed of transplant options (% of 2a)	91.2	92.6	92.9	93.3	93.7	93.7	93.5	
2u 2u	Patients not informed of transplant options (n)	1	92.0 1	1	95.5 1	1.0	1.0	1.0	
2u 2v	Reason not informed (% of 2u; may not sum to 100%) ^{*B}	1	1	1	1	1.0	1.0	1.0	
2.V	Medically unfit	39.2	44.4	43.7	22.5	25.6	25.9	25.2	
	Unsuitable due to age	0.0	44.4 0.0	43.7	0.1	0.1	0.1	0.1	
	Psychologically unfit	0.0	0.0	0.1	0.1	0.0	0.0	0.1	
	Patient declined information	0.0 7.4	11.8	0.0 7.5	4.5	3.6	3.5	3.2	
	Patient declined information Patient has not been assessed	40.0	46.9	43.5	4.5	33.2	33.6	34.9	
	Patient has not been assessed	40.0	40.9	43.5	23.9	55.2	55.0	54.9	
	Comorbid Conditions								
2w	Pre-existing comorbidity (% yes of 2a)	•							
	Congestive heart failure	26.2	25.4	25.7	27.9	28.1	28.1	27.7	
	Atherosclerotic heart disease	10.6	10.2	10.7	19.6	19.5	19.5	19.3	
	Other cardiac disorder	20.3	19.7	20.9	25.6	25.7	25.7	25.3	
	CVD, CVA, TIA	8.4	8.2	8.2	9.9	10.0	10.0	9.8	
	Peripheral vascular disease	7.7	7.0	7.5	9.6	9.6	9.7	9.6	
	History of hypertension	86.7	86.6	87.3	88.5	89.1	89.2	89.1	
	Diabetes	63.8	62.7	62.8	63.1	63.5	63.5	63.7	
	Diabetes on insulin	41.8	40.5	40.1	40.4	41.1	41.2	41.0	
	COPD	8.5	7.9	7.7	8.6	8.6	8.6	8.2	
	Current smoker	6.9	6.9	7.0	8.2	8.1	8.1	7.7	
	Cancer	6.7	6.7	6.8	8.5	8.5	8.5	8.3	
	Alcohol dependence	1.9	1.9	1.9	2.4	2.4	2.4	2.3	
	Drug dependence	1.6	1.7	1.7	2.4	2.5	2.5	2.5	
	Inability to ambulate	6.5	5.9	5.9	9.4	9.3	9.3	9.3	
	Inability to transfer	3.4	3.1	3.0	5.5	5.6	5.6	5.7	
2x	Average number of comorbid conditions	3.0	2.9	3.0	3.3	3.3	3.3	3.3	
	Standardized Modality Switch Ratio (SMoSR)	2021	2022	2023	2021-2023		2021-2023 *A		
2y	Eligible patients (n)	14	14	13	41 *C	15.0	15.0	14.9	
-, 2z	Patient-years at risk (n)	10.1	9.6	9.5	29.2 ^{*C}	10.6	10.6	10.5	
2aa	Number of modality switches (n)	1	1	1	4*C	1.3	1.3	1.3	
2ab	Expected number of modality switches (n)	1.4	1.4	1.4	4.0 ^{*C}	1.3	1.3	1.3	
2ac	SMoSR (2aa/2ab) *D				1.47	1.05	1.05	1.00	
2ad	P-value for SMoSR *E				0.503	n/a	n/a	n/a	
2ae	Confidence interval for SMoSR ^{*E}								
	High (97.5% limit)				8.51	n/a	n/a	n/a	
	Low (2.5% limit)				0.31	n/a	n/a	n/a	

n/a = not applicable

[*A - *F] See Appendix: Table Footnotes on page 29.

[*1] See Guide, Section V for detailed information regarding the summaries included in Table 2.

[*2] For continuous variables, summaries include only values in range indicated in brackets.

SAMPLE Dialysis Facility State: XX Network: 99 CCN: SAMPLE

TABLE 3: Mortality Summary for All Dialysis Patients (2021-24) & New Dialysis Patients (2021-23) *1

				This Fa	cility		Regional Averages *A				
	Measure Name	2021	2022	2023	2024	2021-2024	State	Network	U.S.		
	All Patients: Death Counts							2021-2024			
3a	Patients (n)	83	82	82	81	328 * ^C	82.3	82.3	90.3		
3b	Patient-years at risk (n)	59.2	58.2	58.1	58.1	233.6 ^{*C}	58.5	58.5	59.9		
3c	Deaths (n)	12	11	10	10	43 * ^C	10.8	10.8	11.5		
3d	Expected deaths (n)	13.1	11.9	10.9	11.0	44.8 ^{*C}	11.2	11.2	11.5		
3e	Withdrawal from dialysis prior to death (% of 3c)	20.9	22.2	23.1	23.4	22.3	22.7	22.6	21.8		
3f	Death due to Infections (% of 3c)	10.3	9.9	9.5	9.2	9.8	9.7	9.7	9.6		
	Death due to Cardiac causes (% of 3c)	43.8	45.4	46.2	46.1	45.1	45.6	45.7	44.7		
	Death due to Liver disease (% of 3c)	1.2	1.3	1.3	1.3	1.2	1.2	1.3	1.2		
3g	Dialysis unrelated deaths (n; excluded from SMR)	0	0	0	0	0 *C	0.1	0.1	0.1		
	All Patients: Standardized Mortality Ratio (SMR)										
3h	SMR (3c/3d) *D, *2	0.95	0.98	0.99	1.00	0.98	0.97	0.97	1.00		
3i	P-value for SMR ^{*E}	0.498	0.497	0.498	0.499	0.497	n/a	n/a	n/a		
3j	Confidence interval for SMR ^{*E}										
	High (97.5% limit)	1.88	1.89	1.92	1.95	1.58	n/a	n/a	n/a		
	Low (2.5% limit)	0.50	0.50	0.48	0.48	0.63	n/a	n/a	n/a		
3k	SMR percentiles for this facility										
	In this State	50	50	50	50	50	n/a	n/a	n/a		
	In this Network	50	50	50	50	50	n/a	n/a	n/a		
	In the U.S.	50	50	50	50	50	n/a	n/a	n/a		
	New Patients: First-Year Death Counts	2021	2022	2023		2021-2023		2021-2023			
31	New patients (n)	15	15	15		45 ^{*C}	15.4	15.4	15.3		
3m	Patient-years at risk (n)	13.7	13.3	13.3		40.3 ^{*C}	13.7	13.7	13.7		
3n	Deaths (n)	3	2	2		8 *C	2.6	2.6	2.6		
30	Expected deaths (n)	2.8	2.5	2.5		7.7 ^{*C}	2.5	2.5	2.5		
3p	Withdrawal from dialysis prior to death (% of 3n)	22.6	22.8	24.2		22.8	23.3	23.2	23.0		
3q	Death due to Infections (% of 3n)	9.3	9.2	9.2		9.2	8.9	9.0	9.0		
	Death due to Cardiac causes (% of 3n)	40.6	40.3	40.2		39.9	39.9	40.0	40.3		
	Death due to Liver disease (% of 3n)	2.3	2.4	2.0		2.3	2.2	2.2	2.2		
	First-Year Standardized Mortality Ratio (FySMR)										
3r	FySMR (3n/3o) *D, *2	0.95	0.95	0.95		1.01	1.01	1.01	1.00		
3s	P-value for FySMR ^{*E}	0.496	0.501	0.502		0.497	n/a	n/a	n/a		
3t	Confidence interval for FySMR *E										
	High (97.5% limit)	2.49	2.52	2.57		2.20	n/a	n/a	n/a		
	Low (2.5% limit)	0.35	0.35	0.33		0.42	n/a	n/a	n/a		
3u	FySMR percentiles for this facility										
	In this State	50	50	51		51	n/a	n/a	n/a		
	In this Network	50	50	51		51	n/a	n/a	n/a		
	In the U.S.	49	50	51		51	n/a	n/a	n/a		

n/a = not applicable

[*A - *F] See Appendix: Table Footnotes on page 29.

[*1] See *Guide, Section VI* for detailed information regarding the summaries included in Table 3.

[*2] The SMRs (3h, 3r) are not shown if there are fewer than 3 expected deaths.

TABLE 4: Hospitalization Summary for Medicare Dialysis Patients, 2021-2024^{*1}

				This Fa	cility		Reg	ional Averag per Year	es ^{*A} ,
	Measure Name	2021	2022	2023	2024	2021-2024	State	Network	U.S.
	Medicare Dialysis Patients							2021-2024	
4a	Medicare dialysis r differences (n)	67	65	64	64	260 ^{*C}	65.2	65.2	70.9
4b	Patient-years at risk (n)	45.8	44.7	44.6	44.4	179.5 ^{*C}	45.0	45.0	45.7
	Standardized Hospitalization Ratio (SHR) for								
	Days Hospitlizated *2								
4c	Total days hospitalized (n)	508	509	499	498	2,014 ^{*C}	505.0	504.9	543.3
4d	Expected total days hospitalized (n)	559.9	558.1	553.0	559.4	2,125.4 ^{*C}	531.4	531.3	543.3
4e	SHR (Days) $(4c/4d)^{*D}$	0.95	0.95	0.95	0.95	0.95	0.95	0.95	1.00
4f	P-value for SHR (Days) ^{*E}	0.503	0.503	0.503	0.499	0.500	n/a	n/a	n/a
4g	Confidence interval for SHR (Days) *E								
-	High (97.5% limit)	1.97	1.97	1.94	1.90	1.73	n/a	n/a	n/a
	Low (2.5% limit)	0.56	0.56	0.57	0.57	0.64	n/a	n/a	n/a
4h	SHR (Days) percentiles for this facility								
	In this State	50	50	50	50	49	n/a	n/a	n/a
	In this Network	50	50	50	50	49	n/a	n/a	n/a
	In the U.S.	50	50	50	50	49	n/a	n/a	n/a
	Standardized Hospitalization Ratio (SHR) for Hospital Admissions *2, *3				V				
4i	Total admissions (n)	62	59	63	64	248 *C	62.2	62.2	64.6
4j	Expected total admissions (n)	65.6	62.7	67.3	70.2	253.1 ^{*C}	63.3	63.3	64.6
4k	SHR (Admissions) (4i/4j) *D	0.99	0.98	0.98	0.98	0.98	0.98	0.98	1.00
41	P-value for SHR (Admissions) ^{*E}	0.500	0,498	0.501	0.501	0.499	n/a	n/a	n/a
4m	Confidence interval for SHR (Admissions) ^{*E}								
	High (97.5% limit)	1.72	1.72	1.68	1.69	1.53	n/a	n/a	n/a
	Low (2.5% limit)	0.61	0.62	0.63	0.62	0.69	n/a	n/a	n/a
4n	SHR (Admissions) percentiles for this facility								
	In this State	50	50	50	50	49	n/a	n/a	n/a
	In this Network	50	50	50	50	49	n/a	n/a	n/a
	In the U.S.	50	50	50	50	49	n/a	n/a	n/a
40	Diagnoses associated with hospitalization (% of 4a)								
	Septicemia	12.3	12.2	12.2	12.3	12.3	12.7	12.6	12.2
	Acute myocardial infarction	7.2	7.4	7.8	8.2	7.6	7.9	7.9	7.5
	Congestive heart failure	27.7	28.4	29.2	29.9	28.8	29.6	29.6	28.3
	Cardiac dysrhythmia	18.9	18.8	19.3	19.7	19.2	19.6	19.6	18.6
	Cardiac arrest	2.9	2.8	2.5	2.5	2.7	2.8	2.8	2.7
4p	One day admissions (% of 4i)	8.8	8.4	8.4	8.1	8.4	8.3	8.2	8.2
4q	Average length of stay (days per admission; 4c/4i)	7.8	8.1	7.8	7.6	7.8	7.8	7.8	8.1
4r	Admissions that originate in the ED (% of 4i)	80.5	81.5	81.5	81.7	81.3	83.7	84.0	84.2

SAMPLE Dialysis Facility State: XX Network: 99 CCN: SAMPLE

TABLE 4 (cont.): Hospitalization Summary for Medicare Dialysis Patients, 2021-2024^{*1}

ardized Emergency Department Encounter Ratio R) ^{*2} gency department (ED) events (n)	2021	2022	2023	2024	2021-2024	-	Regional Averages ^{*A} , per Year				
R) ^{*2}					4041-404 4	State	Network	U.S.			
ency department (ED) events (n)											
	36	30	27	25	117 ^{*C}	29.5	29.5	29.5			
ted number of emergency department events (n)	37.5	32.2	29.0	26.9	119.6 ^{*C}	29.9	29.9	29.9			
(4s/4t) *D	1.04	1.04	1.05	1.05	1.02	0.99	0.99	1.00			
e for SEDR [*] E	0.505	0.504	0.502	0.503	0.499	n/a	n/a	n/a			
lence interval for SEDR *E											
High (97.5% limit)	2.15	2.17	2.18	2.21	1.89	n/a	n/a	n/a			
Low (2.5% limit)	0.53	0.53	0.53	0.53	0.58	n/a	n/a	n/a			
ardized Ratio of ED Encounters Occurring											
30 Days of Hospital Discharge (ED30) *2											
discharges (n)	30	25	23	22	100 *C	25.4	25.4	25.4			
ED visits within 30 days of hospital discharge (n)	4	4	3	3	14 * ^c	3.6	3.6	3.6			
ted total ED visits within 30 days of hospital discharge (n)	5.1	4.5	4.2	3.9	16.7 ^{*C}	4.2	4.2	4.2			
Ratio $(4y/4z)^{*E}$	1.06	1.07	1.07	1.07	1.04	0.92	0.92	0.91			
e for ED30 Ratio ^{*E}	0.499	0.498	0.495	0.494	0.500	n/a	n/a	n/a			
lence interval for ED30 Ratio ^{*E}											
High (97.5% limit)	2.46	2.54	2.60	2.64	1.94	n/a	n/a	n/a			
Low (2.5% limit)	0.34	0.32	0.32	0.31	0.47	n/a	n/a	n/a			
ardized Readmission Ratio (SRR) *2			•				2024				
discharges (n)	57	56	57	58		62.3	62.3	62.4			
readmissions (n)	15	14	15	16		16.9	16.9	17.0			
ted total readmissions (n)	16	15	16	17		17.2	17.3	17.3			
4ae/4af) *D	0.99	1.00	1.00	0.99		1.02	1.03	1.03			
e for SRR ^{*E}	0.501	0.499	0.498	0.498		n/a	n/a	n/a			
dence interval for SRR ^{*E}	-										
High (97.5% limit)	1.63	1.65	1.63	1.60		n/a	n/a	n/a			
Low (2.5% limit)	0.53	0.54	0.54	0.55		n/a	n/a	n/a			
ie le H	ae/4af) *D for SRR *E nce interval for SRR *E ligh (97.5% limit) ow (2.5% limit)	ae/4af) *D 0,99 for SRR *E 0.501 nce interval for SRR *E 1.63 ligh (97.5% limit) 0.53	ae/4af) *D 0,99 1.00 for SRR *E 0.501 0.499 nce interval for SRR *E 1.63 1.65	ae/4af) *D 0.99 1.00 1.00 for SRR *E 0.501 0.499 0.498 nce interval for SRR *E 1.63 1.65 1.63 jow (2.5% limit) 0.53 0.54 0.54	ae/4af) *D 0.99 1.00 1.00 0.99 for SRR *E 0.501 0.499 0.498 0.498 nce interval for SRR *E 1.63 1.65 1.63 1.60 igh (97.5% limit) 0.53 0.54 0.54 0.55	ae/4af) *D 0.99 1.00 1.00 0.99 for SRR *E 0.501 0.499 0.498 0.498 nce interval for SRR *E 1.63 1.65 1.63 1.60 ow (2.5% limit) 0.53 0.54 0.55	ae/4af) *D 0.99 1.00 1.00 0.99 1.02 for SRR *E 0.501 0.499 0.498 0.498 n/a nce interval for SRR *E 1.63 1.65 1.63 1.60 n/a igh (97.5% limit) 0.53 0.54 0.54 0.55 n/a	ae/4af) *D 0.99 1.00 1.00 0.99 1.02 1.03 for SRR *E 0.501 0.499 0.498 0.498 n/a n/a nce interval for SRR *E 1.63 1.65 1.63 1.60 n/a n/a			

n/a = not applicable

[*A - *F] See Appendix: Table Footnotes on page 29.

[*1] See *Guide, Section VII* for detailed information regarding the summaries included in Table 4. SEDR and ED30 statistics include only patients with Medicare as primary insurer and exclude patients with Medicare Advantage plans. All other summaries include patients with Medicare as primary insurer or with Medicare Advantage plan.

[*2] SHRs and SEDR are not shown if there are less than five patient-years at risk. SRR and ED30 ratios are not shown if fewer than 11 index discharges in the year.

[*3] Rows 4i-4n exclude index COVID hospitalizations, while rows 4p-4r include all admissions. Row 40 includes diagnoses in any position on a hospital inpatient claim.

TABLE 5: Transplantation Summary for Dialysis Patients under Age 75, 2021-2024

				This Fa	cility		Regional Averages ^{*A} , per Year, 2021-2024			
	Measure Name	2021	2022	2023	2024	2021-2024	State	Network	U.S	
	All Transplants									
5a	Eligible patients (n)	67	65	64	64	260 ^{*C}	65.1	65.1	71.7	
5b	Transplants (n)	2	2	3	3	10 *C	2.5	2.5	2.5	
5c	Donor type (sums to 5b) *B									
	Living donor (n)	0	0	0	0	2^{*C}	0.5	0.5	0.5	
	Deceased donor (n)	2	2	2	2	8 *C	2.0	2.0	2.1	
	First Transplants									
5d	Eligible patients (n)	62	61	60	60	243 *C	60.9	61.0	66.7	
5e	Patient-years at risk (n)	44.4	43.4	43.0	42.7	173.5 *C	43.5	43.5	44.5	
5f	First transplants (n) *2	2	2	2	2	● 9 ^{*C}	2.2	2.2	2.2	
5g	Expected first transplants (n)	2.1	2.2	2.3	2.4	8.7 ^{*C}	2.1	2.1	2.2	
	Standardized 1st Transplantation Ratio (STR) *3									
5h	STR (5f/5g) *D					1.04	1.05	1.04	1.00	
5i	P-value for STR *E					0.390	n/a	n/a	n/a	
5j	Confidence interval for STR *E									
	High (97.5% limit)					2.02	n/a	n/a	n/a	
	Low (2.5% limit)					0.50	n/a	n/a	n/a	
5k	STR percentiles for this facility									
	In this State					50	n/a	n/a	n/a	
	In this Network					50	n/a	n/a	n/a	
	In the U.S.			Ť		50	n/a	n/a	n/a	

n/a = not applicable

[*A - *F] See Appendix: Table Footnotes on page 29.

[*1] See Guide, Section VIII for detailed information regarding the summaries included in Table 5.

[*2] Among first transplants that occurred after the start of dialysis from 2021-2024, 3.2% of transplants in the U.S. were not included because the transplant occurred fewer than 91 days after the start of ESRD and 0.7% were not included because the patient was not assigned to a facility at time of transplant.

[*3] This section is calculated for the 4-year period only and not reported if there are fewer than 3 expected transplants.

SAMPLE Dialysis Facility State: XX Network: 99 CCN: SAMPLE

TABLE 6: Waitlist Summary for All Dialysis Patients (2021-2024) & New Dialysis Patients (2021-2023) *1

				This Fa	cility		Reg	gional Averag	es ^{*A}
	Measure Name	2021	2022	2023	2024	2021-2024	State	Network	U.S.
	All Dialysis Patients under Age 75							2024	
6a	Eligible patients (n)	66	64	63	62		62.0	62.1	62.1
6b	Patient-months at risk (n) *F	558	543	537	530		531.6	531.8	531.9
6c	Patient-months on the waitlist (% of 6b) *F	16.6	16.4	16.1	16.2		16.3	16.1	16.5
6d	Patient-months on the waitlist by subgroup (%) *F, *2								
	Age < 40	24.9	24.2	24.1	24.7		25.5	25.5	25.8
	Age 40-74	15.5	15.4	15.1	15.1		15.2	15.1	15.5
	Male	17.7	17.5	17.1	17.4		17.4	17.2	17.7
	Female	15.1	14.8	14.5	14.5		14.6	14.5	14.7
	African American	16.9	16.7	16.3	16.2		15.5	15.4	15.3
	Asian/Pacific Islander	23.9	24.0	23.6	22.8		24.2	23.6	24.0
	Native American	12.1	11.8	10.9	12.7		13.5	13.4	9.6
	White, Hispanic	17.3	16.4	16.2	16.5		15.2	15.0	16.7
	White, non-Hispanic	16.4	16.0	15.7	16.0		16.6	16.5	16.5
	Other/unknown race	15.0	15.0	14.9	14.4		15.0	14.8	15.7
	Diabetes	13.8	13.9	13.7	13.8		13.9	13.7	14.1
	Non-diabetes	18.8	18.4	17.9	17.9		18.1	18.1	18.5
	Previous kidney transplant	29.0	28.0	27.3	26.8	•	27.7	27.8	28.2
	No previous kidney transplant	15.7	15.6	15.3	15.4		15.4	15.3	15.7
	< 2 years since start of ESRD	12.3	12.7	12.7	13.1		13.8	13.7	13.6
	2-4 years since start of ESRD	21.2	21.0	20.8	20.7		20.8	20.7	21.1
	5+ years since start of ESRD	17.5	16.6	15.9	15.6		14.7	14.6	15.5
6e	Age-adjusted percentage of patient-months waitlisted *3	16,7	16.5	16.1	16.2		16.2	16.2	16.2
6f	P-value for percent waitlisted *E	0.488	0.487	0.493	0.495		n/a	n/a	n/a
6g	Confidence interval for percent waitlisted *E		· ·						
	High (97.5% limit)	43.7	43.5	42.9	42.8		n/a	n/a	n/a
	Low (2.5% limit)	6.0	5.8	5.7	5.7		n/a	n/a	n/a
	Dialysis Patients in Active Status on the Waitlist	*							
6h	Patient-months on the waitlist in active status (% of 6b) $^{*_{\rm F}}$	10.2	9.6	9.4	9.5		9.6	9.5	9.8
6i	Age-adjusted percentage of patient-months	10.3	9.7	9.4	9.6		9.5	9.5	9.6
	waitlisted in active status ^{*3}								
6j	P-value for percent waitlisted in active status*E	0.509	0.513	0.521	0.519		n/a	n/a	n/a
6k	Confidence interval for percent waitlisted in active status *E								
	High (97.5% limit)	35.6	34.7	34.1	34.0		n/a	n/a	n/a
	Low (2.5% limit)	3.1	2.8	2.8	2.8		n/a	n/a	n/a

SAMPLE Dialysis Facility State: XX Network: 99 CCN: SAMPLE

TABLE 6 (cont.): Waitlist Summary for All Dialysis Patients (2021-2024) & New Dialysis Patients (2021-2023) *1

				This Fa	cility		Reg	ional Average	es ^{*A}
	Measure Name	2021	2022	2023	2024	2021-2024	State	Network	U.S.
	All Patients: Prevelent Standardized Waitlist Ratio (PSWR)							2021-2024 *A	
61	Eligible patients (n)	53	50	48	46	94 ^{*C}	23.6	23.6	23.6
6m	Patient-years at risk (n)	38.9	35.7	34.1	32.9	141.7 ^{*C}	35.6	35.6	35.6
6n	Waitlist events (n) *4	3	3	3	3	11 *C	2.7	2.7	2.7
60	Expected waitlist events (n) *4	2.7	2.7	2.8	3.0	10.7 * ^C	2.7	2.7	2.7
6р	PSWR (6n/60) *D	1.10	1.09	1.11	1.11	1.06	1.04	1.03	1.01
6q	P-value for PSWR *E	0.500	0.499	0.494	0.496	0.490	n/a	n/a	n/a
6r	Confidence interval for PSWR *E								
	High (97.5% limit)	15.40	19.41	96.21	13.46	2.55	n/a	n/a	n/a
	Low (2.5% limit)	0.24	0.25	0.26	0.26	0.36	n/a	n/a	n/a
	New Patients: First-Year Standardized Waitlist Ratio (FySWR)	2021	2022	2023		2021-2023		2021-2023 *A	
6s	Eligible patients (n)	10	10	10		▲ 30 *C	10.3	10.3	10.3
6t	Patient-years at risk (n)	9.2	8.9	8.7		26.8 ^{*C}	9.2	9.2	9.2
6u	First waitlist events (n) *4	1	1	1		3 *C	0.9	0.9	0.9
бv	Expected first waitlist events (n) ^{*4}	0.9	0.9	1.1		2.8 ^{*C}	0.9	0.9	0.9
бw	FySWR (6u/6v) ^{*D}		•			0.98	1.04	1.03	1.00
6x	P-value for FySWR ^{*E}					0.494	n/a	n/a	n/a
6у	Confidence interval for FySWR ^{*E}								
-	High (97.5% limit)		< _			2.65	n/a	n/a	n/a
	Low (2.5% limit)			7		0.32	n/a	n/a	n/a

n/a = not applicable

[*A - *F] See Appendix: Table Footnotes on page 29.

[*1] See *Guide, Section IX* for detailed information regarding the summaries included in Table 6. Eligible patient-months (6b) include patients assigned to the facility on the last day of each month.

[*2] The waitlist percentage for each subgroup is calculated as a rate of waitlisted patient-months to patient-months at risk in each subgroup. A missing value indicates that there were no eligible patients in the subgroup.

[*3] Age-adjusted percentages of prevalent patients waitlisted (6e,6i) is not shown if there are fewer than 11 eligible patients in this facility.

[*4] An event is defined as a waitlisting or living-donor transplant. FySWR and PSWR are not shown if a facility has less than 2 expected waitlisted events or less than 11 eligible patients.

SAMPLE Dialysis Facility State: XX Network: 99 CCN: SAMPLE

TABLE 7: Influenza Vaccination Summary for All Dialysis Patients, Flu Seasons during August2021 - December 2024*1

			This I	Facility		Regional Averages *A			
	Measure Name	2021	2022	2023	2024	State	Network	U.S.	
							2024		
7a	Eligible patients on 12/31 (n)	62	61	61	61	62.3	62.3	62.4	
7b	Patients excluded due to medical contraindication (n)	1	2	1	1	1.2	1.2	1.2	
	Full Flu Season (Aug. 1-Mar. 31 of following year)						2023		
7c	Patients vaccinated (% of 7a)	82.2	81.2	78.9		79.7	79.8	79.9	
/d	P-value *E (for 7c compared to U.S. value *2)	0.178	0.162	0.161		n/a	n/a	n/a	
/e	Reason for no vaccination (% of 7a)								
	Declined vaccination	8.4	7.6	8.5		8.8	8.7	8.5	
	Other reason or vaccine data not available	8.7	11.2	12.6		11.5	11.5	11.5	
	Half Flu Season (Aug. 1-Dec. 31)						2024		
ſf	Patients vaccinated (% of 7a)	81.0	80.1	77.7	77.0	77.7	77.8	78.0	
7g	P-value ^{*E} (for 7f compared to U.S. value ^{*3})	0.175	0.160	0.159	0.169	n/a	n/a	n/a	
7h	Patients vaccinated by subgroup (%) *4								
	Medicare	81,9	81.2	78.9	78.4	78.8	78.9	79.	
	Medicare Advantage	83.2	82.5	80.1	79.5	79.9	80.0	80.2	
	Medicare as primary insurer	81.2	80.2	77.9	77.2	77.5	77.5	77.	
	Non-Medicare	74.5	74.1	70.9	70.5	71.6	71.8	72.	
	Dual Medicare/Medicaid eligible	80.2	80.0	78.0	76.9	77.3	77.3	77.0	
	Age < 18	69.6	64.3	60.5	64.7	62.4	63.7	62.9	
	Age 18-39	70.9	70.9	67.9	66.9	66.7	66.8	67.	
	Age 40-64	79.1	78.3	74.9	74.2	74.4	74.5	75.0	
	Age 65-74	83.6	82.8	80.8	79.9	80.3	80.3	80.	
	Age 75+	85.6	85.1	83.6	83.3	83.9	84.0	83.9	
	Male	80.8	79.7	77.0	76.3	76.8	76.9	77.	
	Female	81.3	81.1	79.0	78.5	78.8	78.8	78.	
	African American	77.7	77.9	74.7	74.2	73.6	73.4	74.	
	Asian/Pacific Islander	85.0	84.9	82.7	83.3	84.3	84.5	84.:	
	Native American	82.3	80.3	78.9	78.9	78.3	79.2	82.	
	White	82.1	81.0	78.8	77.9	78.8	79.0	79.	
	Other/unknown race	76.2	75.5	73.3	74.3	75.2	75.2	74.	
	Hispanic	82.9	82.2	80.2	79.9	81.0	81.4	82.	
	< 1 year since start of ESRD	73.9	71.5	70.8	70.3	70.2	70.2	70.	
	1-2 years since start of ESRD	81.8	80.8	77.1	76.4	76.5	76.5	76.	
	3+ years since start of ESRD	83.8	84.1	81.4	80.7	81.4	81.5	81.	

n/a = not applicable

[*A - *F] See Appendix: Table Footnotes on page 29.

[*1] See *Guide*, Section X for detailed information regarding the summaries included in Table 7. All Dialysis Patients are those treated on December 31st of the reporting year.

[*2] Compared to the U.S. value for that year and time period (8/1-3/31): 82.5% (2021), 82.2% (2022), 79.9% (2023).

[*3] Compared to the U.S. value for that year and time period (8/1-12/31): 81.3% (2021), 81.1% (2022), 78.7% (2023), 78.0% (2024).

[*4] A missing value indicates that there were no eligible patients in the subgroup.

TABLE 8: Anemia Management Summaries for Adult Dialysis Patients, 2021-2024^{*1}

			This l	Facility		Region	al Averages*	ges ^{*A} , 2024	
	Measure Name	2021	2022	2023	2024	State	Network	U.S.	
	Hemoglobin and ESA for Adult Hemodialysis (HD) Pati	ents							
8a	Eligible HD patients (n)	69	68	67	67	74.0	74.1	74.1	
8b	Eligible HD patient-months (n) *F	600	587	586	584	644.1	645.0	645.0	
8c	Average hemoglobin (g/dL) (average of 8b)	10.7	10.7	10.7	10.7	10.7	10.7	10.7	
8d	Hemoglobin categories (% of 8b; sums to 100%)								
	<10 g/dL	23.2	23.6	23.2	23.5	23.4	23.4	23.3	
	10-<11 g/dL	33.2	33.2	32.2	32.5	33.1	33.2	33.2	
	11-12 g/dL	27.7	27.7	29.0	29.0	29.6	29.6	29.6	
	>12 g/dL	12.3	12.0	12.8	12.7	12.6	12.6	12.6	
	Missing/Out of range	3.5	3.5	2.8	2.3	1.3	1.2	1.3	
8e	ESA prescribed (% of 8b)	75.6	74.9	75.7	76.4	77.4	77.4	77.6	
	Hemoglobin and ESA for Adult Peritoneal Dialysis (PD) Patients			•				
8f	Eligible PD patients (n)	10	10	11	10	24.2	23.8	23.0	
8g	Eligible PD patient-months (n) *F	81	81	82	83	192.1	189.5	188.	
8h	Average hemoglobin (g/dL) (average of 8g)	10.9	10.9	11.0	11.0	11.0	11.0	11.0	
8i	Hemoglobin categories (% of 8g; sums to 100%)								
	<10 g/dL	21.9	22.7	22.4	22.3	22.4	22.4	22.0	
	10-<11 g/dL	27.9	27.6	26.6	27.3	27.9	27.9	28.0	
	11-12 g/dL	25.9	25.5	26.1	26.2	26.7	26.8	26.7	
	>12 g/dL	19.6	19.5	20.6	20.6	20.5	20.5	20.2	
	Missing/Out of range	4.8	4.7	4.2	3.5	2.4	2.4	2.5	
8j	ESA prescribed (% of 8g)	54.0	53.4	53.5	53.5	54.4	54.4	55.1	
	Standardized Transfusion Ratio (STrR)								
8k	Eligible adult Medicare patients (n)	34	29	26	24	26.0	26.0	26.1	
81	Patient-years at risk (n)	20.1	16.9	14.9	13.4	14.5	14.5	14.5	
8m	Total transfusions (n)	7	5	5	4	4.5	4.5	4.5	
8n	Expected total transfusions (n)	7.4	5.9	5.2	4.7	4.7	4.7	4.7	
80	STrR (8m/8n) *D, *E	0.96	0.95	0.95	0.93	0.97	0.97	1.00	
8p	P-value for STrR ^{*E}	0.502	0.498	0.502	0.497	n/a	n/a	n/a	
8q	Confidence interval for STrR ^{*E}								
	High (97.5% limit)	3.33	3.45	3.54	3.50	n/a	n/a	n/a	
	Low (2.5% limit)	0.42	0.41	0.40	0.40	n/a	n/a	n/a	

n/a = not applicable

[*A - *F] See Appendix: Table Footnotes on page 29.

[*1] See *Guide*, *Section XI* for detailed information regarding the summaries included in Table 8. Transfusion summaries include adult Medicare dialysis patients only and not shown if there are fewer than 10 patient-years at risk (81).

SAMPLE Dialysis Facility State: XX Network: 99 CCN: SAMPLE

TABLE 9: Dialysis Adequacy Summaries for Adult Dialysis Patients, 2021-2024^{*1}

			This F	facility		Regional Averages *A , 202			
	Measure Name	2021	2022	2023	2024	State	Network	U.S.	
	Hemodialysis (HD) Adequacy								
9a	Eligible adult HD patients (n)	69	68	67	67	74.0	74.1	74.1	
9b	Eligible adult HD patient-months (n) *F	600	587	586	584	644.1	645.0	645.6	
9c	Average serum albumin (g/dL) (average of 9b)	3.9	3.9	3.9	3.9	3.9	3.9	3.9	
9d	Serum albumin categories (% of 9b; sums to 100%)		• • •	• • •			•		
	< 3.0 g/dL	2.2	1.9	1.8	1.7	1.8	1.8	1.7	
	3.0-<3.5 g/dL	8.9	8.2	8.0	8.0	8.1	8.1	7.9	
	3.5-<4.0 g/dL	36.4	36.8	39.4	39.8	40.6	40.6	40.2	
	>=4.0 g/dL	47.9	48.2	47.1	47.4	47.6	47.6	48.0	
	Missing	4.6	4.8	3.8	3.1	2.0	2.0	2.1	
9e	Serum albumin<4.0 g/dl (% of 9b)	47.5	46.9	49.1	49.5	50.4	50.4	49.9	
9f	Ultrafiltration rate average (ml/kg/hr) (average of 9b)	7.6	7.6	7.5	75	7.6	7.6	7.6	
)g	Ultrafiltration rate categories (% of 9b; sums to 100%)	7.0	7.0	7.5	1.5	7.0	7.0	7.0	
'g	<=13 ml/kg/hr	85.3	84.8	85.4	86.4	88.0	88.0	87.9	
	-	7.5		7.4	7.1	7.4	7.3		
	>13 ml/kg/hr		7.6		7.1	r		7.5	
	Out of Range (=20 ml/kg/hr)	2.3	2.4	2.6	4.0	2.4	2.4	2.3	
.1	Missing	4.9	5.2	4.7	· · · · · · · · · · · · · · · · · · ·	2.2	2.2	2.3	
h.	Eligible adult HD Kt/V patients (n)	65	64	63	63	71.4	71.4	71.:	
)i	Eligible adult HD Kt/V patient-months (n) *F	561	549	547	543	616.0	616.7	617.	
Эj	Average Kt/V ^{*4} (average of 9i) ^{*2}	1.6	1.6	1.6	1.6	1.6	1.6	1.6	
9k	Kt/V categories (% of 9i; sums to 100%) *2		V						
	<1.2	2.3	2.7	2.5	2.4	1.6	1.6	1.6	
	1.2-<1.8	70.6	71.6	69.4	71.3	72.2	72.2	72.0	
	>=1.8	24.3	22.8	25.6	24.1	25.0	25.0	25.	
	Missing/Out of range	2.9	2.9	2.5	2.2	1.2	1.2	1.3	
	Peritoneal Dialysis (PD) Adequacy								
) 1	Eligible adult PD patients (n)	10	10	11	10	24.2	23.8	23.0	
m	Eligible adult PD patient-months (n)*F	81	81	82	83	192.1	189.5	188.	
n	Average weekly Kt/V (average of 9m) *2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	
90	Weekly Kt/V categories (% of 9m; sums to 100%) *2								
	<1.7	6.1	6.2	5.7	5.6	4.8	4.7	4.8	
	1.7-<2.5	65.8	65.4	65.1	65.6	68.0	68.0	68.	
	>=2.5	23.5	23.5	24.3	24.5	24.3	24.4	24.	
	Missing/Out of range	4.6	4.8	4.9	4.2	2.9	2.8	2.9	
)p	Average serum albumin (g/dL) (average of 9m)	3.7	3.7	3.7	3.7	3.7	3.7	3.7	
pq	Serum albumin categories (% of 9m; sums to 100%)								
•	< 3.0 g/dL	6.6	5.9	5.0	5.0	5.1	5.1	4.9	
	3.0-<3.5 g/dL	21.6	20.7	19.8	20.2	20.8	20.8	20.	
	3.5-<4.0 g/dL	40.5	42.1	43.8	44.1	45.0	44.9	45.0	
	>=4.0 g/dL	26.7	26.6	27.1	26.8	26.5	26.5	26.	
	Missing	4.7	4.7	4.4	3.8	20.5	2.6	2.7	
	111001116	· -- ./	/	7.7	5.0	2.0	2.0	2.1	

n/a = not applicable

[*A - *F] See Appendix: Table Footnotes on page 29.

[*1] See Guide, Section XII for detailed information regarding the summaries included in Table 9.

[*2] Kt/V summaries are supplemented with Medicare claims if missing in EQRS. HD Kt/V summaries are restricted to patients who dialyze thrice weekly. See section of Guide titled "Determination of Thrice Weekly Dialysis" for more information. The most recent value over a 4-month period is selected for PD Kt/V.

TABLE 10: Mineral Metabolism Summaries for All Adult Dialysis Patients, 2021-2024^{*1}

			This F	acility		Region	al Averages [*]	^A , 202
	Measure Name	2021	2022	2023	2024	State	Network	U.S.
l0a	Eligible adult patients (n)	78	77	77	77	82.1	82.2	82.4
0b	Eligible adult patient-months (n) *F	683	670	670	668	717.2	718.1	719.
0c	Average Phosphorus (mg/dL) *2	5.5	5.5	5.5	5.6	5.6	5.6	5.6
0d	Phosphorus categories (% of 10b; sums to 100%) *2							
	<3.5 mg/dL	6.7	7.2	7.2	6.9	7.1	7.1	7.1
	3.5-4.5 mg/dL	21.7	22.3	22.1	21.4	21.6	21.6	21.
	4.6-5.5 mg/dL	28.8	28.0	27.2	26.8	27.0	27.1	27.
	5.6-7.0 mg/dL	22.2	22.2	23.5	24.5	24.8	24.8	24.
	>7.0 mg/dL	16.2	15.9	16.3	17.3	17.5	17.5	17.
	Out of Range (<0.01 mg/dL or >20 mg/dL)	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	Missing	4.5	4.4	3.7	3.1	2.0	1.9	2.0
0e	Average calcium uncorrected (mg/dL) (average of 10b)	8.9	8.9	8.9	8.9	8.9	8.9	8.9
0f	Calcium uncorrected categories (% of 10b; sums to 100%)							
	<8.4 mg/dL	18.2	18.6	20.0	19.0	19.2	19.2	19.
	8.4-10.2 mg/dL	76.4	76.2	75.5	76.9	77.9	77.9	77.
	>10.2 mg/dL	1.3	1.2	1.1	1.3	1.2	1.2	1.
	Out of Range (<0.01 mg/dL or >20 mg/dL)	0.0	0.0	0.0	0.0	0.0	0.0	0.
	Missing	4.1	4.0	3.5	2.9	1.7	1.7	1.
)g	Average uncorrected serum or plasma calcium > 10.2 mg/dL (%) *3	3.0	2.8	2.4	2.1	1.2	1.2	1.

n/a = not applicable

[*A - *F] See Appendix: Table Footnotes on page 29.

[*1] See Guide, Section XIII for detailed information regarding the summaries included in Table 10.

[*2] Eligible patients included in the phosphorus summaries differ slightly from what is reported in 10b since it includes patient-months within the first 90 days of ESRD.

[*3] Hypercalcemia is averaged from uncorrected serum or plasma calcium values over a rolling 3-month period. Eligible patients included in the hypercalcemia summary differs slightly from what is reported in 10b since patients must be 18 years old as of the first day of the 3-month period. Missing or out of range values a counted towards the numerator.

TABLE 11: Vascular Access Information for All Dialysis Patients and Access-Related Infection Summaries for All Medicare Patients, 2021-2024^{*1}

			This I	Facility		Regio	nal Averages*	[^] , 2024
	Measure Name	2021	2022	2023	2024	State	Network	U.S.
	Vascular Access							
11a	Prevalent adult hemodialysis patients (n)	73	72	71	70	77.6	77.7	77.7
11b	Prevalent adult hemodialysis patient-months (n) *F, *2	626	612	606	603	665.3	666.3	666.9
11c	Vascular access type in use (% of 11b; sums to 100%)							
	Arteriovenous fistula	59.4	58.2	57.8	57.2	58.0	57.9	58.6
	Arteriovenous graft	16.3	15.9	16.0	16.3	17.0	17.1	16.7
	Catheter	20.8	22.3	23.3	24.2	23.9	23.9	23.6
	Other/Missing	3.6	3.5	2.8	2.2	1.1	1.1	1.1
11d	Standardized Fistula Rate (SFR) ^{*D, *3}	60.5	59.4	58.9	58.2	58.1	58.2	58.1
11e	P-value for SFR *E	0.512	0.512	0.513	0.509	n/a	n/a	n/a
11f	Confidence interval for SFR *E							
	High (97.5% limit)	35.5	34.6	34.1	32.8	n/a	n/a	n/a
	Low (2.5% limit)	82.4	81.8	81.6	81.5	n/a	n/a	n/a
11g	Long-Term Catheter Rate *4	15.9	17.4	18.2	18.7	18.0	18.0	17.9
	Vascular Access at First Treatment							
11h	Incident hemodialysis patients (n)	14	13	13	13	14.7	14.8	14.
11i	Vascular access type in use (% of 11h; sums to 100%)							
	Arteriovenous fistula	13.2	13.1	13.5	13.0	12.2	12.2	12.
	Arteriovenous graft	2.9	3.2	3.6	3.6	3.4	3.4	3.4
	Catheter	82.1	81.7	81.0	79.4	80.6	80.6	80.
	Other/Missing	1.9	1.9	1.9	4.0	3.8	3.7	3.8
11j	Arteriovenous fistulae in place (% of 11h) *5	13.8	13.7	14.3	13.5	12.7	12.7	12.
	Infection: Peritoneal dialysis (PD)							
11k	Eligible PD patients (n)	6	5	5	5	9.6	9.4	9.4
111	Eligible PD patient-months *F	40	36	33	31	66.2	65.0	64.
11m	PD catheter infection rate per 100 PD patient-months	2.72	2.68	2.78	2.64	2.43	2.43	2.4
11n	P-value *E of 11m (compared to U.S. value)*6	0.378	0.399	0.391	0.410	n/a	n/a	n/a

n/a = not applicable

[*A - *F] See Appendix: Table Footnotes on page 29.

[*1] See Guide, Section XIV for detailed information regarding the summaries included in Table 11.

[*2] Patient months with a catheter that have limited life expectancy, including under hospice care in the current reporting month, or with metastatic cancer, end stage liver disease, coma or anoxic brain injury in the past 12 months, were excluded.

[*3] Includes patients with an autogenous arteriovenous (AV) fistula as the sole means of vascular access. SFR is calculated as an adjusted rate of AV fistula in use reported in 11c; not shown if fewer than 11 eligible adult HD patients.

[*4] Includes patients using a catheter continuously for three months or longer. Patients with other or missing access types (11c) are also counted as catheter in use in the numerator.

[*5] Includes all patients with fistulae, regardless of whether or not they received their hemodialysis treatments using their fistulae.

[*6] Compared to U.S. value for that year: 2.59 (2021), 2.47 (2022), 2.47 (2023), and 2.41 (2024).

TABLE 12: Comorbidities Reported on Inpatient Medicare Claims for Medicare Dialysis Patients Treated as of December 31st of Each Year, 2021-2024^{*1}

			This l	Facility		Region	al Averages*	A , 202
	Measure Name	2021	2022	2023	2024	State	Network	U.S
12a	Medicare dialysis patients on 12/31 (n)	46	45	45	45	47.8	47.9	48.0
12b	Comorbidity (% yes of 12a)							
	Infections							
	AIDS/HIV positive	0.7	0.7	0.7	0.7	0.8	0.8	0.8
	Intravascular/implanted device-related *2	4.8	4.9	4.9	4.7	4.6	4.6	4.7
	Hepatitis B	0.4	0.4	0.4	0.4	0.4	0.4	0.4
	Hepatitis other	1.6	1.5	1.4	1.3	1.3	1.3	1.4
	Metastatic	1.4	1.5	1.6	1.5	1.6	1.6	1.0
	Pneumonia	3.2	3.4	3.7	3.7	3.7	3.7	3.
	Tuberculosis	0.2	0.2	0.3	0.2	0.3	0.3	0.
	Other	18.0	18.5	18.8	18.8	18.7	18.6	19
	Cardiovascular							
	Cardiac arrest	1.0	1.0	0.9	0.9	0.9	0.9	0.
	Cardiac dysrhythmia	18.8	19.0	19.7	19.8	19.9	19.8	19
	Cerebrovascular disease	7.2	7.3	7.6	7.7	7.9	7.9	8.
	Congestive heart failure	30.0	30.8	31.9	32.1	32.4	32.3	32
	Ischemic heart disease	25.8	26.2	26.6	26.5	26.7	26.7	26
	Myocardial infarction	7.4	7.9	8.3	8.5	8.6	8.6	8.
	Peripheral vascular disease *3	16.7	16.7	17.4	17.2	17.5	17.5	17
	Other							
	Alcohol dependence	1.3	1.4	1.5	1.4	1.4	1.4	1.
	Anemia	2.9	3.2	3.3	3.3	3.3	3.3	3.
	Cancer	3,4	3.5	3.7	3.8	3.8	3.8	3.
	Chronic obstructive pulmonary disease	14.3	14.6	14.9	15.0	15.1	15.0	14
	Diabetes	35.9	36.4	37.0	36.4	36.7	36.7	37
	Drug dependence	1.2	1.2	1.3	1.2	1.2	1.2	1.
	Gastrointestinal tract bleeding	3.0	3.0	3.1	3.2	3.3	3.3	3.
	Hyperparathyroidism	20.0	20.7	21.8	22.2	22.4	22.2	22
12c	Average number of comorbid conditions	2.2	2.2	2.3	2.3	2.3	2.3	2.

n/a = not applicable

[*A - *F] See Appendix: Table Footnotes on page 29.

[*1] See *Guide, Section XV* for detailed information regarding the summaries included in Table 12. Based on patients with Medicare as primary insurer on 12/31 each year.

[*2] This category includes bloodstream and other infections related to intravascular access and other implanted devices, not limited to dialysis access.

[*3] Peripheral vascular disease includes venous, arterial and nonspecific peripheral vascular diseases.

TABLE 13: Facility Information, 2024

		This Facility	Regi	onal Averag 2024	ges ^{*A} ,
	Measure Name	2024	State	Network	U.S.
13a	Organization	SAMPLE MEDICAL CARE(SMC)			
13b	Ownership	Profit			
13c	Initial Medicare certification date	01/01/2016			
13d	Number of stations	17			
13e	Services provided	Unavailable			
13f	Shifts after 5:00 pm	Yes			
13g	Dialyzer Reuse	Yes			
13h	CMS Certification Numbers (CCN) included in this report	999999			
13i	National Provider Identifier (NPI)	1234569874			
	Long Term Care (LTC)				
13j	Dialysis facility located in a Skilled Nursing Facility (SNF)	Yes			
13k	Services provided in LTC facility by non-SNF based facility	None			
	Patient Placement				
131	Patients treated during year from AFS Form-2744 (n)	103	102.6	102.7	102.9
13m	Transferred into facility (% of 131)	18.2	18.2	18.2	17.9
13n	Transferred out of facility (% of 13l)	18.9	18.7	18.7	18.5
13o	Patients treated on 12/31 (n)	67	n/a	n/a	n/a
13p	Medicare eligibility status (% of 13o; sums to 100% ^{*B})		n/a	n/a	n/a
-	Medicare	80.8	80.9	80.9	80.1
	Medicare application pending	1.2	1.1	1.1	1.1
	Non-Medicare	17.9	18.0	18.0	18.8
	Survey and Certification *2				
13q	Date of last survey	01/03/2018			
13r	Type of survey	Unknown			
13s	Compliance condition after survey	Meets Requirements			
13t	Number of CFC deficiencies cited	0	0.2	0.2	0.2
13u	Number of Standard deficiencies cited	4	4.2	4.2	4.2

n/a = not applicable

[*A - *F] See Appendix: Table Footnotes on page 29.

[*1] See *Guide, Section XVI* for detailed information regarding the summaries included in Table 13 Information based on data reported in EQRS as of May 2025. If missing, data were not available.

[*2] Data on this section are from the facility's latest survey since January 2009 according to information reported in iQIES as of early June 2025.

SAMPLE Dialysis Facility State: XX Network: 99 CCN: SAMPLE

TABLE 14: Selected Measures for Dialysis Patients under Age 18, 2021-2024 *1

			This I	facility		Region	al Averages [*]	^A , 2024
	Measure Name	2021	2022	2023	2024	State	Network	U.S.
	Patient Characteristics							
14.1a	Patients treated on 12/31 (n)	7	7	7	6	n/a	n/a	n/a
14.1c	Age (% of 14.1a; sums to 100%)							
	< 5	25.2	23.9	23.4	25.7	28.8	29.5	29.8
	5-9	14.2	11.4	10.8	11.0	11.8	11.9	12.6
	10-14	30.0	35.6	33.0	27.3	26.5	26.0	26.3
	15-17	30.6	29.0	32.7	36.0	33.0	32.5	31.3
14.1d	Female (% of 14.1a)	43.8	42.1	39.8	40.7	39.7	39.7	41.3
14.1e	Race (% of 14.11a; sums to 100%)							
	African American	26.6	24.8	27.9	26.6	28.7	29.0	26.3
	Asian/Pacific Islander	7.1	7.2	5.7	6.3	4.1	4.3	4.7
	Native American	0.5	1.2	1.5	2.1	1.2	1.1	1.2
	White	64.0	63.8	59,3	<u>5</u> 3.7	55.1	54.7	57.1
	Other/Unknown/Missing	1.8	3.0	5.6	11.3	11.0	10.9	10.6
14.1f	Ethnicity (% of 14.1a; sums to 100%)							
	Hispanic	31.3	30.6	30.6	30.4	32.1	32.3	35.1
	Non-Hispanic	68.6	69.3	69.4	69.5	67.5	67.3	64.7
	Unknown	0.0	0.0	0.0	0.1	0.4	0.4	0.2
14.1g	Cause of ESRD (% of 14.1a; sums to 100%)							
	Diabetes	0.3	0.3	0.2	0.1	0.3	0.3	0.3
	Hypertension	1.1	2.1	1.6	2.1	2.0	1.9	1.6
	Glomerulonephritis	32.5	31.0	27.1	27.2	27.7	27.4	26.6
	Cystic Kidney	31,3	34.4	37.1	34.9	35.9	37.0	37.0
	Congenital/Hereditary	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	Hemolytic Uremic Syndrome	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	Other	28.0	26.4	28.4	28.3	24.9	24.6	25.0
	Unknown/Missing	6.8	5.9	5.6	7.4	9.3	8.8	9.6
14.1i	Years since start of ESRD (% of 14.1a; sums to 100%)							
	<1	34.8	32.7	35.5	34.0	35.1	34.7	33.8
	1-2	21.8	24.9	23.8	23.1	23.8	24.5	24.4
	2-3	12.2	10.6	12.2	12.8	11.2	11.0	11.3
	3-6	16.6	15.3	13.6	13.7	15.0	15.1	15.4
	6+	14.6	16.5	14.9	16.5	14.9	14.7	15.1
l4.1k	Modality (% of 14.1a; sums to 100%)							
	In-center hemodialysis	46.4	47.4	52.4	55.5	54.0	53.6	53.6
	Home hemodialysis	0.6	0.9	0.4	0.1	0.7	0.7	0.5
	Continuous ambulatory peritoneal dialysis	1.7	2.0	2.2	1.8	0.9	1.0	1.1
	Continuous cycling peritoneal dialysis	51.2	49.3	44.4	42.4	44.2	44.5	44.6
	Other modality	0.1	0.3	0.6	0.1	0.3	0.3	0.3

SAMPLE Dialysis Facility State: XX Network: 99 CCN: SAMPLE

TABLE 14 (cont.): Selected Measures for Dialysis Patients under Age 18, 2021-2024 *1

			This H	Facility		Region	al Averages [*]	^A , 2024
	Measure Name	2021	2022	2023	2024	State	Network	U.S.
	Characteristics of New Dialysis Patients							
14.2a	Total number of patients with forms (n)	4	4	4	4	n/a	n/a	n/a
14.2g	Medical coverage (% of 14.2a; sums to 100%)							
U	Employer group only	20.7	19.3	14.7	18.4	18.1	18.3	18.7
	Medicare (alone or combined w/ other insurance)	0.7	1.1	2.1	1.9	1.6	1.6	1.6
	Medicaid only	59.5	61.9	61.0	56.4	53.7	53.2	52.9
	Other/Unknown/None	19.0	17.8	22.3	23.3	26.6	26.9	26.8
14.2k	Number of incident hemodialysis patients (n)	2	2	2	2	n/a	n/a	n/a
14.21	Access used at first outpatient dialysis (% of 14.2k; sums to 100	%)						
	Arteriovenous fistula	1.3	3.6	0.8	1.5	1.4	1.3	1.2
	Arteriovenous graft	1.0	0.3	0.0	1.0	0.2	0.2	0.2
	Catheter	97.3	96.0	98.7	65.4	68.6	67.0	66.5
	Other/Unknown/Missing	0.4	0.0	0.5	32.1	29.8	31.5	32.1
14.2m	Arteriovenous fistulae placed (% of 14.2k)	2.8	5.0	1.6	2.2	2.3	1.9	1.9
14.2s	Pre-ESRD nephrologist care (% of 14.2a; sums to 100%)							
	No	21.8	20.5	21.6	15.9	19.0	18.7	18.7
	Yes, < 6 months	23.0	23.6	23.2	22.6	25.6	25.8	26.1
	Yes, 6-12 months	16.3	15.5	15.0	12.4	11.7	11.6	11.6
	Yes, > 12 months	33.2	34.3	34.0	39.9	33.5	33.6	34.0
	Unknown	5.6	6.2	6.2	9.1	10.2	10.3	9.7
14.2t	Informed of transplant options (% of 14.2a)	91.9	93.5	93.8	66.2	64.9	64.7	63.6
	Death Rates	\sim	•					
14.3a	Patients (n)	11	11	12	11	n/a	n/a	n/a
14.3c	Deaths (n)	0	0	0	0	n/a	n/a	n/a
	Hospitalization Statistics							
14.4a	Medicare dialysis patients (n)	4	3	3	3	n/a	n/a	n/a
14.4c	Total days hospitalized (n)	33	32	35	29	n/a	n/a	n/a
14.4i	Total admissions (n)	4	4	4	4	n/a	n/a	n/a
	Transplantation							
14.5d	Eligible patients (n)	10	10	10	10	n/a	n/a	n/a
14.5f	First transplants (n)	3	3	3	3	n/a	n/a	n/a
	Waitlist							
14.6b	Eligible patients-months (n) *F	92	94	94	91	n/a	n/a	n/a
14.6c	Patients-months on the waitlist (% of 14.6b)	34.9	33.8	34.0	35.6	32.6	32.7	31.6
14.6d	Patient-months on the waitlist by age (%)							
	Age < 10	33.6	31.1	30.1	31.2	29.4	29.3	27.8
	Age 10-17	42.4	42.2	39.4	39.2	36.6	36.1	36.1

SAMPLE Dialysis Facility State: XX Network: 99 CCN: SAMPLE

TABLE 14 (cont.): Selected Measures for Dialysis Patients under Age 18, 2021-2024^{*1}

			This H	Facility		Region	al Averages*	^A , 2024
	Measure Name	2021	2022	2023	2024	State	Network	U.S.
	Hemoglobin							
14.8b	Eligible HD patient-months (n) *F	34	37	38	39	n/a	n/a	n/a
14.8c	Average hemoglobin (g/dL) (average of 14.8b)	10.7	10.8	10.8	10.8	10.8	10.8	10.8
14.8d	Hemoglobin categories (% of 14.8b; sums to 100%)	1017	1010	1010	1010	1010	1010	1010
1 1100	<10 g/dL	22.6	21.9	22.7	21.3	20.2	20.5	19.6
	10-<11 g/dL	27.1	26.3	22.4	23.6	23.4	23.9	22.8
	11-12 g/dL	31.1	29.6	30.8	30.4	29.9	30.6	30.1
	> 12 g/dL	14.1	14.6	15.7	16.5	14.7	14.6	14.5
	Missing/Out of Range	5.1	7.5	8.4	8.2	11.7	10.3	13.0
14.8g	Eligible PD patient-months (n) *F	42	42	41	37	n/a	n/a	n/a
14.8h	Average hemoglobin (g/dL) (average of 14.8g)	10.9	11.0	10.8	10.8	10.8	10.8	10.9
14.8i	Hemoglobin categories (% of 14.8g; sums to 100%)	1000	1110	1010		1010	1010	100
	<10 g/dL	24.6	21.9	25.7	26.4	26.7	26.7	26.0
	10-<11 g/dL	23.5	22.8	22.7	23.5	22.3	23.1	22.8
	11-12 g/dL	21.9	22.3	21.5	22.0	22.5	22.7	23.3
	> 12 g/dL	19.1	19.5	17.8	19.6	19.3	18.9	19.7
	Missing/Out of Range	10.8	13.5	12.3	8.4	9.2	8.5	8.3
	Albumin							
14.9b	Eligible HD patient-months (n) *F	34	37	38	39	n/a	n/a	n/a
14.9c	Average serum albumin (g/dL) (average of 14.9b)	4.1	4.1	4.0	4.0	4.0	4.0	4.0
14.9d	Serum albumin categories (% of 14.9b; sums to 100%)							
	< 3.0 g/dL	2.3	2.1	2.5	2.3	2.2	2.4	2.4
	3.0-<3.5 g/dL	6.1	6.0	5.8	6.9	6.9	6.9	7.6
	3.5-<4.0 g/dL	26.0	22.7	25.1	27.1	27.2	27.4	27.8
	>=4.0 g/dL	59.9	61.4	57.9	55.2	51.6	52.8	48.9
	Missing	5.6	7.9	8.8	8.5	12.0	10.5	13.2
14.91	Eligible PD patient-months (n) *F	42	42	41	37	n/a	n/a	n/a
14.90	Average serum albumin (g/dL) (average of 14.91)	3.7	3.7	3.7	3.7	3.7	3.7	3.7
14.9p	Serum albumin categories (% of 14.91; sums to 100%)							
	< 3.0 g/dL	9.5	9.2	8.0	8.6	9.3	9.2	10.7
	3.0-<3.5 g/dL	15.5	15.5	14.9	17.4	17.3	17.6	17.8
	3.5-<4.0 g/dL	26.5	26.0	27.6	27.3	28.1	28.9	27.8
	>=4.0 g/dL	36.6	33.9	32.8	30.4	29.0	29.2	29.0
	Missing	11.9	15.4	16.7	16.3	16.3	15.1	14.8
	<i>Kt/V</i> *2							
14.9h	Eligible HD patient-months (n) *F	26	30	31	32	n/a	n/a	n/a
14.9j	HD: $Kt/V \ge 1.2$ (% of 14.9h)	90.6	88.0	87.5	86.4	84.0	86.2	82.6
14.9n	PD: Kt/V >= 1.8 (% of 14.9l)	77.2	74.1	77.0	75.6	77.7	78.1	79.0
	Vascular Access *3							
14.11b	Eligible patient-months (n) *F	39	42	43	44	n/a	n/a	n/a
14.11c	Arteriovenous Fistula in use (%)	17.4	42 14.5	45 11.1	11.1	9.8	10.0	9.1
14.11c	Long-Term Catheter Rate	57.5	64.1	66.0	67.9	70.2	70.8	72.5

n/a = not applicable

[*A - *F] See Appendix: Table Footnotes on page 29.

[*1] See *Guide, Section XVII* for detailed information regarding the summaries included in Table 14.

[*2] Kt/V: K = dialyzer clearance of urea; t = dialysis time; V = patient's total body water. HD Kt/v summaries restricted to patients on thrice weekly in-center hemodialysis. PD Kt/V summaries select the most recent value collected within 6 months of the reporting month.

[*3] 'Arteriovenous Fistula in use' includes patients with an autogenous arteriovenous (AV) fistula as the sole means of vascular access. 'Long-Term Catheter Rate' includes patients using a catheter continuously for three months or longer. Patients with other or missing access types (11c) are also counted as catheter in the numerator.

TABLE 15: Selected Measures for Nursing Home (NH) Dialysis Patients, 2021-2024 *1

			This F	facility		Region	al Averages [*]	A, 202
	Measure Name	2021	2022	2023	2024	State	Network	U.S
	Patient Characteristics							
15.1a	Patients treated on 12/31 (n)	14	14	15	17	n/a	n/a	n/a
15.1c	Age (% of 15.1a; sums to 100%)							
	18-64	33.8	33.2	32.4	32.2	32.1	32.1	32.
	65+	66.2	66.8	67.6	67.8	67.9	67.9	67.
15.1d	Female (% of 15.1a)	48.2	48.1	48.0	47.0	46.6	46.7	46.
15.1e	Race (% of 15.1a; sums to 100%)							
	African American	35.0	36.0	36.4	36.3	35.5	35.3	37.
	Asian/Pacific Islander	4.9	4.9	5.1	5.2	5.2	5.4	4.6
	Native American	1.0	1.1	1.0	1.1	1.2	1.2	1.2
	White	58.7	57.7	56.9	56.3	57.0	57.0	56.
	Other/Unknown/Missing	0.4	0.4	0.6	1.0	1.1	1.1	1.0
15.1f	Ethnicity (% of 15.1a; sums to 100%)							
	Hispanic	16.2	16.3	16.5	16.8	17.2	17.0	15.
	Non-Hispanic	83.7	83.7	83.5	83.2	82.7	82.9	84.
	Unknown	0.0	0.0	0.0	0.1	0.1	0.1	0.
15.1g	Cause of ESRD (% of 15.1a; sums to 100%)	•						
	Diabetes	56.7	54.8	54.0	53.0	52.5	52.4	51.
	Hypertension	26.5	27.6	27.9	28.3	28.3	28.2	28.
	Glomerulonephritis	4.6	4.4	4.7	4.4	4.3	4.3	4.3
	Other	12.1	13.0	13.2	14.0	14.6	14.6	14.
	Unknown/Missing	0.2	0.2	0.2	0.4	0.4	0.4	0.4
15.1i	Years since start of ESRD (% of 15.1a; sums to 100%)							
	<1	21.8	20.9	20.3	20.4	20.5	20.6	20.
	1-2	14.8	16.0	15.2	15.1	15.4	15.3	15.
	2-3	12.5	11.6	12.9	12.2	12.2	12.2	12.
	3-6	25.6	25.3	24.9	25.6	25.6	25.6	25.
	6+	25.4	26.2	26.7	26.7	26.3	26.3	26.
15.1k	Modality (% of 15.1a; sums to 100%)							
	In-center hemodialysis	94.4	94.0	93.7	93.2	91.2	91.1	90.
	Home hemodialysis	2.6	3.3	3.8	4.4	6.1	6.1	7.
	Continuous ambulatory peritoneal dialysis	0.2	0.1	0.1	0.2	0.2	0.2	0.2
	Continuous cycling peritoneal dialysis	2.0	1.7	1.6	1.6	1.9	1.8	1.9
	Other modality	0.8	0.9	0.7	0.6	0.6	0.6	0.7

SAMPLE Dialysis Facility State: XX Network: 99 CCN: SAMPLE

TABLE 15 (cont.): Selected Measures for Nursing Home (NH) Dialysis Patients, 2021-2024^{*1}

			This I	Facility		Region	al Averages*	^A , 202
	Measure Name	2021	2022	2023	2024	State	Network	U.S
	Characteristics of New Dialysis Patients							
15.2a	Total number of patients with forms (n)	6	6	6	6	n/a	n/a	n/a
15.2g	Medical coverage (% of 15.2a; sums to 100%)	0	0	Ũ	Ũ	11/ 4		11/0
	Employer group only	3.2	2.7	2.3	2.7	2.4	2.4	2.4
	Medicare (alone or combined w/ other insurance)	78.2	80.7	81.9	81.0	81.8	81.8	82.
	Medicaid only	12.0	11.0	10.8	11.0	10.4	10.3	9.
	Other/Unknown/None	6.6	5.5	4.9	5.3	5.4	5.4	5.4
15.2k	Number of incident hemodialysis patients (n)	6	6	6	7	n/a	n/a	n/:
15.21	Access used at first outpatient dialysis (% of 15.2k; sums to 100%)	0	0	0	,	n/u	11/ u	11/
	Arteriovenous fistula	6.3	6.5	7.4	6.9	6.5	6.5	6.
	Arteriovenous graft	1.8	2.8	2.9	3.1	2.6	2.6	2.:
	Catheter	89.7	88.7	87.5	87.7	88.2	88.3	88.
	Other/Unknown/Missing	2.2	2.1	2.2	2.3	2.6	2.6	2.
15.2m	Arteriovenous fistulae placed (% of 15.2k)	15.7	15.2	16.5	14.5	13.8	13.7	13.
15.2s	Pre-ESRD nephrologist care (% of 15.2a; sums to 100%)	15.7	10.2	10.5	14.5	15.0	15.7	15
13.25	No	19.2	17.5	16.8	16.7	16.9	17.0	16
	Yes, < 6 months	19.2	21.5	21.6	21.8	22.0	22.0	22
	Yes, 6-12 months	17.6	17.3	17.2	17.8	16.0	15.9	15
	Yes, > 12 months	21.4	21.2	21.7	22.2	20.1	20.1	20
	Unknown	21.4	22.4	22.5	21.3	24.8	24.8	20
15.2t	Informed of transplant options (% of 15.2a)	86.3	88.5	88.0	70.5	68.4	68.3	68
	Mortality summary for all NH dialysis patients							
15.3b	Patient-years at risk (n)	14.9	15.3	16.3	17.8	n/a	n/a	n/a
15.3c	Deaths (n)	6	6	7	7	n/a	n/a	n/:
15.3d	Expected deaths (n)	5.8	5.5	5.4	5.8	n/a	n/a	n/:
15.3c/15.3b	Rate (deaths per 100 patient-years at risk)	43.2	42.3	40.0	35.6	39.3	39.3	42
15.3h	SMR (15.3c/15.3d)	1.12	1.15	1.21	1.13	1.24	1.24	1.3
15.3e	Withdrawal from dialysis prior to death (% of 15.3c)	26.3	26.7	27.8	27.9	26.8	26.8	26
15.50		20.5	20.7	27.0	21.9	20.0	20.0	20
	Hospitalization summary for NH Medicare dialysis patients							
15.4b	Patient-years at risk (n)	13.0	13.2	14.1	15.4	n/a	n/a	n/
15.4i	Total admissions (n)	37	36	41	45	n/a	n/a	n/
15.4j	Expected total admissions (n)	24.8	25.0	28.0	30.8	n/a	n/a	n/a
15.4i/15.4b	Rate (admissions per 100 patient-years at risk)	287.0	277.7	288.7	290.3	296.6	296.0	305
5.4k	SHR (15.4i/15.4j)	1.56	1.51	1.50	1.49	1.51	1.51	1.5
5.40	Diagnoses associated with hospitalization (% of 15.4a)							
	Septicemia	25.5	25.0	25.0	24.7	25.5	25.5	22
	Acute myocardial infarction	12.3	12.8	13.5	13.3	13.6	13.6	11
	Congestive heart failure	47.2	48.8	49.9	49.9	49.8	49.8	45
	Cardiac dysrhythmia	35.2	35.0	35.5	35.7	35.8	35.8	32
	Cardiac arrest	4.8	4.4	4.3	4.0	4.3	4.3	3.

SAMPLE Dialysis Facility State: XX Network: 99 CCN: SAMPLE

TABLE 15 (cont.): Selected Measures for Nursing Home (NH) Dialysis Patients, 2021-2024^{*1}

			This I	Facility		Region	al Averages*	^A , 2024
	Measure Name	2021	2022	2023	2024	State	Network	U.S.
	Comorbidities reported on Medicare claims							
15.12a	Medicare dialysis patients on 12/31 (n)	12	12	13	15	n/a	n/a	n/a
15.12b	Comorbidity (% yes of 15.12a)							
	Infections							
	AIDS/HIV positive	0.9	1.0	1.1	1.1	1.0	1.0	1.1
	Intravascular/implanted device-related	8.2	8.4	8.3	7.8	7.8	7.8	8.2
	Hepatitis B	0.9	1.0	0.9	0.8	0.8	0.8	0.8
	Hepatitis other	3.4	3.4	3.1	2.8	2.8	2.7	2.8
	Metastatic	3.6	4.1	4.0	4.2	4.3	4.3	4.6
	Pneumonia	9.2	9.6	9.8	10.2	10.2	10.1	10.6
	Tuberculosis	0.3	0.3	0.3	0.3	0.3	0.3	0.3
	Other	39.6	39.4	40.1	40.0	40.4	40.4	41.2
	Cardiovascular				•			
	Cardiac arrest	2.5	2.3	2.3	2.4	2.3	2.3	2.5
	Cardiac dysrhythmia	38.4	38.8	39.2	39.2	39.4	39.3	39.7
	Cerebrovascular disease	19.1	19.3	19.9	19.8	19.8	19.8	20.1
	Congestive heart failure	54.2	55.7	57.0	57.0	56.9	56.9	57.6
	Ischemic heart disease	47.5	46.8	47.0	46.7	46.5	46.5	46.8
	Myocardial infarction	14.1	14.8	16.0	16.1	16.2	16.2	16.4
	Peripheral vascular disease	36.7	35.6	36.9	36.7	36.5	36.5	36.9
	Anemia	A X						
15.8b	Eligible adult HD patient-months (n) *F	164	168	179	196	n/a	n/a	n/a
15.8d	Hemoglobin (HD) < 10 g/dL (% of 15.8b)	31	31	30	31	30.7	30.6	30.9
15.8e	ESA prescribed (% of 15.8b)	81	80	81	82	81.7	81.6	81.5
15.8g	Eligible adult PD patient-months (n) *F	5	5	5	5	n/a	n/a	n/a
15.8i	Hemoglobin (PD) < 10 g/dL (% of 15.8g)	28	29	28	29	28.3	28.1	28.0
15.8j	ESA prescribed (% of 15.8g)	59	58	58	57	56.6	56.5	55.6
	Dialysis Adequacy *2							
15.9b	Eligible adult HD Kt/V patient-months (n) *F	156	159	170	185	n/a	n/a	n/a
15.9k	Kt/V (HD) < 1.2 (% of 15.9b)	2	3	2	2	2.2	2.2	2.2
15.9m	Eligible adult PD Kt/V patient-months (n) *F	5	5	5	5	n/a	n/a	n/a
15.90	Kt/V (PD) < 1.7 (% of 15.9m)	11	12	10	11	9.1	9.1	8.7
	Vascular Access *3							
15.11b	Prevalent adult hemodialysis patient months (n) *F	171	175	181	196	n/a	n/a	n/a
15.11c	Arteriovenous fistula in use (% of 15.11b)	49.7	48.1	48.8	48.4	48.9	48.9	48.3
15.11g	Long-Term Catheter Rate	21.0	22.5	22.7	22.9	22.8	22.8	23.0

n/a = not applicable

[*A - *F] See Appendix: Table Footnotes on page 29.

[*1] See Guide, Section XVIII for detailed information regarding the summaries included in Table 15.

[*2] Kt/V summaries are supplemented with Medicare claims if missing in EQRS. HD Kt/V summaries are restricted to patients who dialyze thrice weekly. See section of Guide titled "*Determination of Thrice Weekly Dialysis*" for more information. PD Kt/V summaries select the most recent value collected within months of the reporting month.

[*3] Patient months with a catheter that have limited life expectancy, including under hospice care in the current reporting month, or with metastatic cancer, end stage liver disease, coma or anoxic brain injury in the past 12 months, were excluded. Other or missing access types are counted as catheter in use in the numerator. LTCR includes patients using a catheter continuously for three months or longer.

SAMPLE Dialysis Facility State: XX Network: 99 CCN: SAMPLE

Appendix: Table Footnotes

This appendix provides information that applies to multiple tables.

[*A] Regional values are shown for the average facility. When the summary includes three- or four-years, values are shown for the average facility, annualized.

[*B] Values may not sum to exactly 100%. Refer to the appropriate section of the Guide found in the table-specific footnotes (section II) for additional details.

[*C] Sum of counts over the three- or four-years used for calculations and should not be compared to regional averages.

[*D] This value is the ratio of observed events to expected events. This value is not shown when there are too few events to yield reliable result. Refer to the table-specific footnotes (section II) below for the criteria for when the ratio is reported.

[*E] The confidence interval range represents uncertainty in the value of the ratio due to random variation. A p-value less than 0.05 indicates that the difference between the observed and expected events is probably real and is not due to random chance alone. A p-value greater than or equal to 0.05 indicates that the difference could plausibly be due to random chance.

[*F] A patient may be counted up to 12 times per year.

